

Sulphur Iod 3X

Introduction: A mix of sulphur and iodine, this is a remedy often employed on the basis of clinical concepts denoting a tuberculous terrain (Tuberculinum), with recurrent head colds and bronchitis; laryngitis; enlarged adenoids (Agraphis); and a catarrhal condition in the Eustachian tube. The child sleeps with an open mouth.

This is a very profound and long-acting remedy affecting all parts of the body with aggravations morning, afternoon, evening, NIGHT, and after midnight.

Synonym: Iodide of Sulphur

Source: sulphur and iodine

Proved by: Kelsall

Guiding symptoms/Indications:

1. Almost a specific for barber's itch. Cures eczema, when the surface is darkened, swollen, weepy and itches terribly.
2. "pustular eruption on upper lip" and "erythema on chin."
3. Thickness of tongue remaining after parenchymatous glossitis.
-Hypertrophy of parotids after parotitis.
4. Uvula and tonsil enlarged and reddened, accumulation of dark purulent mucus in windpipe, mucus removed with difficulty. In the throat feeling as if swelled.
5. Itching on ears, nose, and in urethra. Papular eruption on the face.
Erythematous breaking out on upper lip, with yellowish pustules, painful, quickly disappearing in dry scales.
6. Boils upon anterior part of neck and in vicinity of the mouth. Arms covered with an itching rash like nettle-rash.

Particulars:

Throat.--Uvula and tonsils enlarged and reddened. Swollen. Tongue thick. Parotid hypertrophied.

Skin.--Itching on ears, nose, and in urethra. Papular eruption on face. Cold-sores on lips. Boils on neck. Barber's itch. Acne. Lichen planus. Arms covered with itching rash. Hair feels as if erect.

Pathogenesis:

Dose and administration: 1 tab 3 to 4 times daily. ½ tab for children.

Side effects: No significant side effect has been observed in proper dosage

Contraindication: There is no known contraindication.

Presentations: 50 tabs (20 gm in PET bottle), 450gm powder in container.