

## Merc Dul 3X

**Introduction:** Catarrhal affections of mucous membranes, especially of the eye and ear. Acute affections of prostate after maltreated stricture. Deafness and catarrhal affections of nares, throat, and pharynx, from mercurial amalgam fillings.

**Synonyms:** Calomel. Subchloride of Mercury. Mercurious chloride. Formula  $Hg_2Cl_2$ .

**Source :** It is a chemical compound of Mercury and Chlorine. Can be purchased from a chemist's shop.

**Preparation :** Triturations are prepared with sugar of milk from which higher potencies are prepared.

**Proved by** Dr. T.F. Allen.

### **Guiding symptoms/Indications:**

1. It is a useful remedy in chronic suppurative otitis media. (Kali m.).
2. Eustachian catarrh with deafness and tinnitus.
3. Salivation with sore mouth and offensive breath. (Bor., Kali phos., Lach., Nit. ac.).
4. Diarrhoea of children, stools grass green (Ipec., Mag. carb.) like chopped eggs (Cham.); profuse; causing soreness of anus.
5. Acute affections of prostate after maltreated stricture. (Acon., Arn., Canth., Ferr. phos.).
6. Ulceration in throat with dysphagia. (Lach., Nit. ac., Sil.).
7. Modalities : worse : night, cold weather  
Better : cold drinks.

### **Particulars:**

Ears.--Otitis media; closure of Eustachian tube; ear troubles of scrofulous children; membrana tympani retracted, thickened and immovable.

Mouth.--Offensive breath; salivation; sore gums. Ulcers. Tongue black. Constant flow of dark, putrid saliva; very offensive. Ulceration of throat, with dysphagia. Granular pharyngitis.

Stomach.--Nausea and vomiting. Cyclic vomiting of infants (Cup ars; Iris).

Stool.--Scanty, bloody mucus, with bile, and constant desire, without tenesmus. Dark-green, watery, with griping. Anus sore and burning. Dysentery; small stools of mucus and blood, covered with bile.

Skin.--Flabby and ill nourished. Swollen glands. Phagedenic ulcers. Copper-colored eruptions.

Relationship.--Compare: Kali mur.

**Dose and administration:** 1 tab 3 to 4 times daily. ½ tab for children.

**Side effects:** No significant side effect has been observed in proper dosage

**Contraindication:** There is no known contraindication.

**Presentations:** 50 tabs (20 gm in PET bottle), 450gm powder in container.